



## King's Research Portal

DOI:

[10.1017/S0007114519000291](https://doi.org/10.1017/S0007114519000291)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Goff, L. M., Huang, P., Silva, M. J., Bordoli, C., Enayat, E. Z., Molaodi, O. R., Cassidy, A., Maynard, M., & Harding, S. (2019). Associations of dietary intake with cardiometabolic risk in a multi-ethnic cohort: a longitudinal analysis of the Determinants of Adolescence, now young Adults, Social well-being and Health (DASH) study. *British Journal of Nutrition*, 121(9), 1069-1079. <https://doi.org/10.1017/S0007114519000291>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

**Associations of dietary intake with cardiometabolic risk in a multi-ethnic cohort: a longitudinal analysis of the Determinants of Adolescence, now young Adults, Social well-being and Health (DASH) study**

Louise M. Goff<sup>1,2</sup>, Peiyuan Huang<sup>1</sup>, Maria J. Silva<sup>1</sup>, Claire Bordoli<sup>1</sup>, Elli Z. Enayat<sup>1</sup>, Oarabile R. Molaodi<sup>3</sup>, Aidan Cassidy<sup>1</sup>, Maria Maynard<sup>4</sup>, Seeromanie Harding<sup>1</sup>

**Affiliations:** <sup>1</sup>Department of Nutritional Sciences, School of Life Course Sciences, Faculty of Life Sciences & Medicine, King's College London, London SE1 9NH <sup>2</sup>Department of Diabetes, School of Life Course Sciences, Faculty of Life Sciences & Medicine, King's College London, London SE1 1UL; <sup>3</sup>MRC/CSO Social and Public Health Sciences Unit, University of Glasgow, Glasgow; <sup>4</sup>School of Clinical & Applied Sciences, Leeds Beckett University, Leeds.

**Pubmed indexing:** Goff, Huang, Silva, Bordoli, Enayat, Molaodi, Cassidy, Maynard, Harding.

**Corresponding author:** Dr Louise M. Goff, Diabetes Research Group, Departments of Diabetes & Nutritional Sciences, School of Life Course Sciences, Faculty of Life Sciences & Medicine, King's College London, Waterloo Campus, Franklin-Wilkins Building, Room 3.87, London, SE1 9NH, United Kingdom. T: +44 (0)20 7848 4273; E: [louise.goff@kcl.ac.uk](mailto:louise.goff@kcl.ac.uk)

**Short running title:** Ethnicity, diet and cardiometabolic risk

**Keywords:** ethnicity, breakfast, diet, cardiometabolic risk, African, Caribbean, South Asian

**Abbreviations:**

BMI: Body Mass Index

CHD: Coronary Heart Disease

DASH: Determinants of Adolescent Social well-being and Health

T2D: type 2 diabetes

WHtR: waist to height ratio

## 1    **ABSTRACT**

2    Unfavourable dietary habits, such as skipping breakfast are common among ethnic minority children  
3    and may contribute to inequalities in cardiometabolic disease. We conducted a longitudinal follow-  
4    up of a subsample of the UK multi-ethnic Determinants of Adolescent Social well-being and Health  
5    (DASH) cohort, which represents the main UK ethnic groups and is now aged 21-23 years. We aimed  
6    to describe longitudinal patterns of dietary intake and investigate their impact on cardiometabolic risk  
7    in young adulthood. Participants completed a dietary behaviours questionnaire and a 24-hr dietary  
8    intake recall; anthropometry, blood pressure, total and HDL-cholesterol and HbA1c were measured.  
9    The cohort consisted of 107 White British, 102 Black Caribbean, 132 Black African, 98 Indian, 111  
10    Bangladeshi/Pakistani and 115 Other/Mixed ethnicity. Unhealthful dietary behaviours such as  
11    skipping breakfast and low intakes of fruit and vegetables were common (56%, 57% and 63%,  
12    respectively). Rates of skipping breakfast and low fruit and vegetable consumption were highest  
13    among Black African and Black Caribbean participants. BMI and cholesterol levels at 21-23 years  
14    were higher among those who regularly skipped breakfast at 11-13 years (BMI 1.41 (95% CI 0.57,  
15    2.26)  $p=0.001$ ; cholesterol 0.15 (95% CI -0.01, 0.31)  $p=0.063$ ) and 21-23 years (BMI 1.05 (95% CI  
16    0.22, 1.89)  $p=0.014$ ; cholesterol 0.22 (95% CI 0.06, 0.37)  $p=0.007$ ). Childhood breakfast skipping is  
17    more common in certain ethnic groups and is associated with cardiometabolic risk factors in young  
18    adulthood. Our findings highlight the importance of targeting interventions to improve dietary  
19    behaviours such as breakfast consumption at specific population groups.

20

21

## 22 INTRODUCTION

23 Ethnic inequalities in a wide range of chronic diseases have been documented <sup>(1)</sup>, with risk factors  
 24 evident early in life <sup>(2)</sup>. Children and youth from ethnic minority groups are more likely to be  
 25 overweight or obese than White-European populations <sup>(3;4)</sup>. The cardiometabolic risks associated with  
 26 obesity are well known, therefore it is not surprising that some ethnic minority populations suffer  
 27 disproportionately high rates of chronic diseases such as type 2 diabetes (T2D), hypertension and  
 28 coronary heart disease (CHD) <sup>(1; 5)</sup>.

29 Ethnic inequalities in obesity and chronic disease risk may be explained by differences in diet and  
 30 lifestyle. Differences in the nutritional composition of the diet are apparent; a recent cohort study of  
 31 children in the UK reported significantly higher energy, fat and sugar intakes among South Asians,  
 32 while Black Africans demonstrated lower intakes of fat and saturated fat but higher intakes of  
 33 carbohydrate, compared to White British <sup>(6)</sup>. Unfavourable dietary habits, such as regularly skipping  
 34 breakfast and consumption of sugar-sweetened fizzy drinks and fast foods, are also more common  
 35 among children and adolescents from some ethnic minority groups <sup>(7; 8; 9; 10)</sup>. The detrimental impact  
 36 of skipping breakfast, and fast food and fizzy drink consumption on childhood and adolescent health  
 37 has been documented through a number of cohort studies, which have shown that those who regularly  
 38 skip breakfast have an increased risk of obesity and detrimental cardiometabolic risk profiles <sup>(10; 11;</sup>  
 39 <sup>12)</sup>. Additionally, studies have shown that children who skip breakfast regularly have poorer quality  
 40 diets, particularly higher energy density, saturated fat and lower micronutrient intakes <sup>(13)</sup>. More  
 41 recently longitudinal studies have investigated the prevalence of these behaviours in young adulthood  
 42 and the detrimental impact of childhood behaviours on adult health <sup>(11; 14; 15)</sup>. Young adulthood is  
 43 recognised as an important transition period, with increasing responsibilities and the development of  
 44 independent health behaviours. The Australian Childhood Determinants of Adult Health (CDAH)  
 45 study found skipping breakfast in both childhood and young adulthood to be associated with larger  
 46 waist circumferences, higher fasting insulin, and higher total- and LDL-cholesterol concentrations  
 47 <sup>(11)</sup>, while in the US-based Project EAT socio-economic status was found to be a principal determinant  
 48 of increases in breakfast skipping between childhood and adulthood <sup>(15)</sup>. To date there have been no  
 49 longitudinal cohort studies to describe ethnic patterns in adolescent and young adult nutritional intake  
 50 and dietary behaviours, and to determine their longitudinal effects on body weight and  
 51 cardiometabolic risk factors. The ‘DASH 10 years on’ study is the first multi-ethnic cohort of young  
 52 adults in the UK and provides longitudinal follow-up data for a subset of the Determinants of  
 53 Adolescent Social well-being and Health (DASH) cohort, who are now young adults (21-23 years).  
 54 We aimed to describe longitudinal patterns of dietary intake in the multi-ethnic Determinants of

55 Adolescent, now young Adults, Social well-being and Health cohort, and investigate the impact of  
56 dietary intake on cardiometabolic risk in young adulthood.

57

## 58    **SUBJECTS AND METHODS**

### 59    *Study Design*

60    Details of the DASH study can be found at <http://dash.sphsu.mrc.ac.uk/> and in a published cohort  
 61    profile <sup>(16)</sup>. The original cohort of 6643 students, aged 11-13y, was recruited between 2002 and 2003,  
 62    from 51 secondary schools in 10 London boroughs and sampled to represent the main ethnic groups  
 63    of the UK population. A pilot follow-up study on 665 participants was completed in March 2014;  
 64    dietary data was collected from 627 participants who took part in the follow-up.

65  
 66    The study was approved by the National Health Service South East Scotland Research Ethics  
 67    Committee (Ref:11/SS/0051). Written informed consent was obtained from participants. The study  
 68    is registered on clinicaltrials.gov ID: NCT03283332.

### 70    *Participants*

71    The subsample consisted of 107 White British, 102 Black Caribbean, 132 Black African, 98 Indian,  
 72    111 Bangladeshi or Pakistani and 115 Other (mainly mixed) ethnicities, and was purposively sampled  
 73    to be representative by gender and socio-economic circumstances (SEC) across the 10 London  
 74    boroughs and 51 schools. Ethnicity was self-declared, measured by self-report using Census  
 75    categories, checked against reported parental ethnicity and grandparents' country of birth.  
 76    Bangladeshis and Pakistanis were approximately equally represented and combined because of  
 77    relatively small sample sizes; participants declaring their ethnicity as White European, White Irish,  
 78    Chinese, mixed ethnicity were classified as 'Other ethnicity'.

### 80    *Procedures*

81    Measurement protocols and questionnaires can be found at <http://dash.sphsu.mrc.ac.uk/>. Nurses and  
 82    research assistants were trained in the assessment methods by the research team for 1 week prior to  
 83    the start of fieldwork, and were re-certified at 6 monthly intervals. All equipment was checked and  
 84    calibrated regularly by the field supervisors. Participants were given a choice of locations (local  
 85    general practitioner's surgeries, local community pharmacies, Clinical Research Centres and Kings  
 86    College London) and assessments took about 2 hours.

### 88    *Dietary behaviours and intake*

89    At age 11-13 and 21-23 years participants completed a short questionnaire regarding their  
 90    consumption of breakfast, fruit and vegetables. For breakfast consumption, participants were given a  
 91    choice of four responses (every day, 3-4 days a week, 1-2 days a week, never/hardly ever). Fruit and

vegetable consumption were separate questions with a choice of seven responses for each (5 or more portions a day, 4 portions a day, 3 portions a day, 2 portions a day, 1 portions a day, I eat fruit/vegetables some days but not every day, I never eat fruit/vegetables). Examples of portions (e.g. handful of grapes, an apple, serving spoon of vegetables, bowl of salad) were given alongside the questions for more accurate estimation of portion size. In addition, at age 11-13 years participants answered an additional question regarding consumption of fizzy drinks, for which a choice of five answers were available (every day, most days, weekly, less than weekly, never).

At 21-23 years dietary intake was assessed using the standardised triple-pass 24-hour recall methodology <sup>(17)</sup>. The study investigator, trained in dietary assessment methods, interviewed the participant face-to-face to recall their food and beverage intake for the preceding 24 hours, extracting details regarding food types, brands, recipes and cooking methods. Detailed information was collected for pre-packaged foods including brands and portion sizes, and full recipes and cooking methods were recorded for home-prepared foods and dishes. Portion sizes were confirmed using a photographic food atlas <sup>(18)</sup> and household measures.

Dietary intake records were coded for analysis by a single coder (CB) and analysed for food and nutrient composition using WISP software, which is based on McCance and Widdowson's *The Composition of Foods*, 6<sup>th</sup> edition. Where food items were not available within the WISP database new entries were created, particularly for home-prepared recipes. Nutritional composition information was sought from food labels and manufacturer websites for branded items, and information for cultural foods were sought from our own database of African and Caribbean dishes <sup>(19)</sup>, the West African Food Table <sup>(20)</sup>, and recipe books. Food weights were assigned from the photo atlases that were used in the interviews. Where food weights were not indicated, average adult portion sizes from the National Diet and Nutrition Survey were used <sup>(21)</sup>.

The quality of reporting in the 24-hour recall records was assessed using a scoring system developed by the researchers. Seven components were identified as important to assess the quality of dietary information (portion sizes, brand information, food description, cooking methods, recipe information, beverage information, intake timings) and these were scored on a scale of 0-2, in which zero indicated no information provided, 1 indicated incomplete information provided, and 2 indicated complete information provided. The scores assigned to each category were totalled and a percentage calculated, indicating the quality of dietary information obtained. For dietary recalls which did not include categories such as composite or homemade dishes these categories were excluded when calculating the diet quality percentage. To test the sensitivity of the protocol to assess the quality of the dietary



information obtained, two researchers trialled a sample of recalls deemed of poor (n=10) and high quality (n=10). Researchers compared quality scores in order to remain consistent. Overall the mean quality score of the dietary records was 69%, ranging from 21 – 100%, with 85% of the recalls scoring  $\geq 50\%$  for the quality of the data recording. In preliminary analysis we investigated the impact of reporting quality on the dietary data to determine whether recalls of poorer quality should be excluded from our final analyses; we found no statistical difference in reported energy intake when records scoring  $< 50\%$  were excluded, therefore, the final analyses included data from the whole cohort.

Macronutrient intakes were expressed as percentage of total energy intake. Dietary fibre (AOAC method) was expressed in grams per 1000 kcals and sodium intake in milligrams per 1000 kcals. Energy intakes were compared to estimated energy requirements to assess potential under-reporting; age and sex specific standard equations were used to estimate basal metabolic rate (BMR)<sup>(22)</sup>. These were multiplied by a lower cut-off physical activity level (PAL) of 1.2 to provide estimates of minimum total energy expenditure and recognise probable under-reporting<sup>(23)</sup>; 36% of participants reported an energy intake below this cut-off and were considered probable under-reporters (rates did not vary by gender, ethnicity or educational attainment (data not shown)). Data from these participants were not excluded from further analyses.

#### ***Body size, biological, and social measures***

Height was measured without shoes, using portable stadiometers, to the nearest 0.1 cm. Weight was measured in light clothing using Salter electronic scales, to the nearest 0.1 kg. Body Mass Index (BMI) was derived as weight (kg)/height (m)<sup>2</sup>; BMI  $> 25$  kg/m<sup>2</sup> was defined as overweight/obese. Waist circumference (cm) was measured midway between the 10<sup>th</sup> rib and the top of the iliac crest, and 0.5 cm subtracted to correct for measurement over T-shirt or vest. The mean of two duplicate measures was derived, with waist to height ratio calculated. Systolic blood pressure (sBP) was measured using validated OMRON M5-1 instruments and appropriately sized cuffs, after sitting quietly for a timed 5 minutes, with  $> 1$  minute between 3 subsequent readings. The mean of the second and third readings was used in analysis. A 25 ml blood sample was taken by venepuncture for the assessment of total and HDL cholesterol and glycated haemoglobin (HbA1c); lipids were analysed using colour photometric assays on an automated clinical chemistry analyser (Siemens Advia 2400 analyser), HbA1c was analysed using HPLC (Premier 9210 analyser). Physical activity was measured using ActivPal<sup>TM</sup> monitors (PAL Technologies, Glasgow UK). The monitor, validated for identifying sitting, standing and walking<sup>(24)</sup> was worn continuously on the front of the thigh for 5 days; the number of minutes spent sitting or in sedentary activity per day was derived and used as the measure of physical inactivity.

162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196

A self-complete questionnaire measured current smoking behaviour and SEC (i.e. educational attainment and employment status).

### *Statistical Analysis*

The dietary behaviours (skipping breakfast, fizzy drink, fruit, and vegetable consumption) were recoded into dichotomous variables as follows: skipping breakfast was defined as not consuming breakfast every day; fizzy drink consumption was defined as consuming fizzy drinks every day or most days; low daily fruit consumption was defined as consuming less than 2 portions of fruit per day; and low daily vegetable consumption was defined as consuming less than 3 portions of vegetables per day. Missing data in each categorical variable were recoded as 'not stated'. Missing data among continuous variables were 0.001% for gender, 0.15% for BMI, 4.5% for waist:height ratio, 25.0% for total cholesterol, 27.2% for HDL-cholesterol, 27.7% for HbA1c, 0.15% for systolic blood pressure, and 5.6% for all six nutritional composition variables.

Continuous variables were tested for normality using the Shapiro-Wilk test, and no transformation was necessary. Ethnic differences in anthropometry, cardiometabolic risks factors and dietary intake were performed using ANOVA with post-hoc analysis. Longitudinal patterns in dietary behaviours (skipping breakfast, fruit and vegetable intake) from 11-13 years to 21-23 years were examined with mixed-effects logistic regression, adjusted for age, gender and ethnicity. Multivariate linear regression was used to examine the association between dietary behaviours at 11-13 years and at 21-23 years and cardiometabolic risk factors at 21-23 years, adjusted for gender and ethnicity at 11-13 years, and additionally adjusted for educational attainment (attainment/non-attainment of higher education degree), physical activity (based on tertiles of sitting/sedentary time), smoking (current smoker/non-smoker), energy intake and saturated fat intake at 21-23 years. N=664, 635, 499, 484, 481, and 664, respectively, for linear regression models with BMI, waist:height ratio, total cholesterol, HDL-cholesterol, HbA1c, and systolic blood pressure, respectively, as the outcome. Multivariate linear regression was used to examine the association between dietary behaviours at 11-13 years and at 21-23 years and nutritional composition of the diet at 21-23 years, adjusted for gender, and ethnicity at 11-13 years, and additionally adjusted for educational attainment (attainment/non-attainment of higher education degree), physical activity (based on tertiles of sitting/sedentary time), and smoking (current smoker/non-smoker) at 21-23 years. N=628 for linear regression models for all six nutritional composition variables as the outcome. All analyses were performed using STATA (version 13; StataCorp LP, College Station, Texas, USA) and statistical significance was considered at  $p<0.05$ .

## 197 RESULTS

198 The cohort consisted of 329 men and 336 women. The representation by ethnicity was: 107 White  
 199 British, 102 Black Caribbean, 132 Black African, 98 Indian, 111 Bangladeshi or Pakistani and 115  
 200 Other (mainly mixed ethnicity). The mean age of the cohort was 22.7 years, mean BMI 24.7 kg/m<sup>2</sup>,  
 201 39% were overweight or obese (BMI>25 kg/m<sup>2</sup>), 24% were smokers and 47% were educated to  
 202 degree level. Rates of smoking were highest among White British (47%) and lowest among Black  
 203 African (11%) participants. Attainment of a university degree was highest among Black African  
 204 (58%) and lowest among Black Caribbean (34%) participants.

205 Dietary data, collected for 627 participants, are shown in Table 1 alongside anthropometry and  
 206 cardiometabolic risk factors for the whole cohort and for each ethnic group at 21-23 years. At 21-23  
 207 years, overall, 56% of participants reported skipping breakfast on a regular basis, 57% reported  
 208 consuming less than 2 portions of fruit a day, and 63% reported consuming less than 3 portions of  
 209 vegetables a day. Ethnic differences in diet, anthropometry and cardiometabolic risk were evident.  
 210 Black Caribbean and Black African participants were significantly heavier than the Indian and  
 211 Pakistani/Bangladeshi participants, and had higher BMI compared with White British and Indian  
 212 participants but there were no differences in waist circumference or waist to height ratio. Total  
 213 cholesterol was lowest in the Black African and highest in the Pakistani/Bangladeshi participants,  
 214 while HDL-cholesterol was highest in the Black Caribbean and Black African participants, and lowest  
 215 in the Indian participants. The highest proportion of breakfast skipping occurred among the Black  
 216 Caribbean (66%) and Black African (64%) groups and was lowest in the Indian group (46%),  
 217 although there were no statistical differences. Saturated fat intakes were lowest among the Black  
 218 African and Indian participants and highest among the Pakistani/Bangladeshi participants. Dietary  
 219 fibre intake was highest in the White British and Indian groups and was significantly higher than the  
 220 Black groups or the Pakistani/Bangladeshi participants. The Pakistani/Bangladeshi participants were  
 221 significantly more likely to have a low daily consumption of vegetables but there were no ethnic  
 222 differences in fruit consumption.

223 We examined longitudinal patterns in the diet behaviours; there were significant increases in the rates  
 224 of skipping breakfast (37 vs 56%  $p=0.035$ ), low consumption of fruit (46 vs 57%  $p=0.024$ ) and of  
 225 vegetables (57 vs 63%  $p=0.021$ ) from 11-13 years to 21-23 years among the whole cohort (Figure 1).  
 226 Age was associated with risk of unhealthful diet behaviours, independent of gender and ethnicity.  
 227 When they were 21-23 years old, individuals were more likely to skip breakfast (OR: 2.65 (95% CI  
 228 2.00, 3.50),  $p<0.001$ ), have low consumption of fruit (OR: 1.39 (95% CI 1.08, 1.79),  $p=0.011$ ), and  
 229 vegetables (OR: 1.28 (95% CI 0.99, 1.66),  $p=0.059$ ) than when they were 11-13 years old, although

the association for low vegetable consumption did not reach statistical significance. After adjusting for gender, ethnicity and educational attainment, skipping breakfast regularly at 11-13 years was a significant determinant of skipping breakfast in young adulthood (OR: 2.36 (95% CI 1.62, 3.44),  $p<0.001$ ). Low fruit consumption and low vegetable consumption at 21-23 years were also significant determinants of skipping breakfast at 21-23 years (Fruit: OR: 2.21 (95% CI 1.54, 3.18),  $p<0.001$ ; Vegetables: OR: 1.91 (95% CI 1.32, 2.76),  $p=0.001$ ) but low fruit consumption and low vegetable consumption at 11-13 years were not significant determinants of skipping breakfast in young adulthood (Fruit: OR: 1.29 (95% CI 0.88, 1.88),  $p=0.190$ ; Vegetables: OR: 1.32 (95% CI 0.91, 1.91),  $p=0.150$ ). Figure 1 shows the rates of (a) skipping breakfast, (b) low fruit and (c) low vegetable consumption at 11-13 years and 21-23 years by ethnic group. Small ethnic sample sizes precluded rigorous testing of ethnic specific effects, nonetheless the descriptive data indicates ethnic variations. Increases in the rates of skipping breakfast were evident in all ethnic groups, with the greatest increase occurring in the White British group. Prevalence of low fruit consumption increased in all ethnic groups, with the Black African group experiencing the greatest increase. While the rates of low vegetable consumption did not increase in either the White British or 'Other' ethnic groups but increased in all other ethnicities, with the greatest increase occurring among the Black African group.

#### *Longitudinal associations between dietary behaviours, anthropometric and cardiometabolic risk factors*

The influence of the dietary behaviours during both adolescence and young adulthood on cardiometabolic risk factors during young adulthood were investigated in multivariate analysis, adjusting for gender, ethnicity, educational attainment, smoking, physical activity, energy and saturated fat intake (Table 2). Skipping breakfast at 11-13 years was a significant determinant of BMI at 21-23 years (1.41 (95% CI 0.57, 2.26),  $p<0.001$ ) as was skipping breakfast at 21-23 years, although the effect was slightly attenuated in this age group (1.05 (95% CI 0.22, 1.89),  $p=0.014$ ). Skipping breakfast at both 11-13 years and 21-23 years were also important determinants of total cholesterol levels (11-13 years: 0.15 (95% CI -0.01, 0.31),  $p=0.063$ ; 21-23 years: 0.22 (95% CI 0.06, 0.37),  $p=0.007$ ). Low intake of fruit at 11-13 years was a significant determinant of waist to height ratio at 21-23 years (0.01 (95% CI 0.00, 0.03),  $p=0.034$ ), and regular fizzy drink consumption at 11-13 years was a significant determinant of systolic blood pressure at 21-23 years (1.77 (95% CI 0.12, 3.43),  $p=0.036$ ).

#### *Longitudinal associations between dietary behaviours and nutritional composition of the diet*

The influence of the dietary behaviours during both adolescence and young adulthood on the nutritional composition of the diet during young adulthood were investigated in multivariate analysis,

263 adjusting for gender, ethnicity, educational attainment, smoking and physical activity (Table 3).  
264 Skipping breakfast at both 11-13 years and 21-23 years were significant determinants of lower dietary  
265 fibre intakes (11-13 years: -0.81 (95% CI -1.49, -0.12),  $p=0.022$ ; 21-23 years: -1.33 (95% CI -2.01, -  
266 0.65),  $p<0.001$ ), as were regular fizzy drink consumption at 11-13 years (-0.72 (95% CI -1.42, -0.02),  
267  $p=0.045$ ), low fruit consumption at 21-23 years (-1.22 (95% CI -1.90, -0.54),  $p<0.001$ ) and low  
268 vegetable consumption at 21-23 years (-1.30 (95% CI -2.04, -0.56),  $p=0.001$ ). Low fruit and vegetable  
269 consumption at 21-23 years were also important determinants of higher saturated fat intakes (fruit:  
270 0.89 (95% CI -0.01, 1.78),  $p=0.052$ ; vegetables: 1.06 (95% CI 0.09, 2.02),  $p=0.032$ ).

271

## 272 DISCUSSION

273 We have undertaken a longitudinal assessment of nutritional intake and dietary behaviours and their  
 274 association with cardiometabolic risk factors among an ethnically diverse cohort of young adults in  
 275 the UK. In this work, we report the detrimental impact of childhood eating behaviours, such as  
 276 regularly skipping breakfast, on cardiometabolic risk factors in young adulthood. We have identified  
 277 an increasing prevalence, from childhood to young adulthood, of less healthy dietary behaviours.  
 278 Furthermore, we have described distinct ethnic patterns, such that skipping breakfast is most prevalent  
 279 among Black African and Caribbean groups.

280 The DASH cohort was recruited primarily to investigate ethnic determinants and patterns of health,  
 281 and through the collection of dietary data we have been able to explore potential predictors of  
 282 cardiometabolic risk in our young adults. In this study, we found the Black African and Caribbean  
 283 groups to have the highest body weights and BMIs, however waist circumference and waist to height  
 284 ratio were comparable to the other groups. In the Indian ethnic group, body weight and BMI were  
 285 significantly lower than the White British and both Black ethnic groups. These findings are in line  
 286 with the existing literature and suggest that higher body weight/BMI amongst African ancestry  
 287 populations is not related to abdominal fat accumulation <sup>(25)</sup>, although fat deposition in the abdominal  
 288 region is usually more prevalent in South Asian populations <sup>(26)</sup>. The Black Africans exhibited low  
 289 total cholesterol and high HDL-cholesterol concentrations; these cardiometabolic risk differences  
 290 may be attributed to dietary differences, namely lower saturated fat, which was also apparent among  
 291 the Black African participants. Among the Indian participants HDL-cholesterol was lower.  
 292 Distinctions in nutritional intake were also evident for this group, namely higher intakes of fibre in  
 293 the Indian group. Ethnic differences in cardiometabolic risk factors have been extensively reported  
 294 and in general our data corroborate the literature. However, our categorisation of ethnicity, which is  
 295 more detailed than that which is often employed, highlights some differences between our findings  
 296 and the existing literature. Higher HbA1c levels amongst African-Americans relative to White-  
 297 Americans have been previously reported <sup>(27; 28)</sup>; in our study the Black Caribbeans and Black  
 298 Africans did not exhibit higher HbA1c levels. It is not clear the extent to which higher HbA1c among  
 299 Black populations is a result of dietary intake versus biological factors. Higher sugar intakes among  
 300 Black Caribbean compared to Black West African adults have previously been described <sup>(19)</sup>, however  
 301 we did not find this in our participants, which is in agreement with Donin *et al.* who reported no  
 302 difference in sugar intake between Black Caribbean and Black African children in the CHASE study  
 303 <sup>(6)</sup>. Lower total cholesterol amongst Black Africans but not Black Caribbeans is in agreement with  
 304 findings from the CHASE study <sup>(2)</sup>, and may relate to lower saturated fat intake amongst Black  
 305 Africans but not Black Caribbeans; a finding that is consistent with that of the CHASE study.

Historically the African-Caribbean diet has been recognised for its low intake of saturated fat, which has been proposed to contribute to the cardioprotective lipid profile that these ethnicities traditionally exhibited <sup>(29; 30; 31)</sup>. However, dietary acculturation and ‘Westernisation’ of the diet results in increasing fat intake, a change that is more apparent amongst Black Caribbean than Black African communities <sup>(19)</sup>, which may explain why the total cholesterol concentration was only lower amongst the Black African group. Lower HDL-cholesterol amongst South Asian populations has been extensively reported in the literature <sup>(31; 32; 33)</sup>, and the nutritional profile that we identified is also consistent with previous reports <sup>(29; 31; 32; 34)</sup>.

Unhealthy dietary behaviours were highly prevalent among our young adults. In agreement with other longitudinal investigations we saw that adolescent breakfast skipping was predictive of the behaviour in adulthood <sup>(11; 14)</sup>. In our study breakfast skipping was particularly prevalent amongst certain ethnic groups. In agreement with other studies of children and young adults <sup>(10; 14; 35)</sup> our Black African and Black Caribbean participants had the highest rates of breakfast skipping, although the White British group saw the greatest increase in skipping breakfast from childhood to young adulthood. Our longitudinal analysis indicated that skipping breakfast in childhood was an important determinant of BMI and total cholesterol concentration in young adulthood. Whilst there have been a number of longitudinal studies to demonstrate the detrimental effects of skipping breakfast on body weight and the development of obesity, there has been very little demonstration of the long-term effects of this behaviour on cardiometabolic risk factors, such as blood lipids. Our study is the first to focus on an ethnically diverse cohort that represents the main ethnic groups of the UK population. Prior to this the long term evidence was limited to the Australian CDAH study, which studied a predominantly White Australian population, and found skipping breakfast in adolescence to have a unfavourable impact on waist circumference, fasting insulin and total- and LDL-cholesterol <sup>(11)</sup>. Furthermore, adolescent and young adult dietary behaviours were seen to be significant determinants of nutritional composition. Skipping breakfast and regular fizzy drink consumption during adolescence were associated with significantly lower intakes of dietary fibre in young adulthood. In young adulthood skipping breakfast and low consumption of fruit and vegetables were associated with significantly lower dietary fibre and significantly higher saturated fat intakes. These findings corroborate with the literature, demonstrating the detrimental impact of skipping breakfast and other dietary behaviours on the quality of the diet <sup>(13)</sup>; these results are particularly alarming given these unhealthful behaviours are most prevalent among some of our ethnic minority groups, who are most at risk of developing obesity and cardiometabolic diseases, such as T2D. Promoting high fibre intakes and restriction of saturated fat are two of the most important public health nutrition campaigns in a bid to reduce chronic disease risk among the general population. Our findings provide a useful insight into dietary

340 behaviours and specific ethnic groups that health promotion campaigns could target in aiming to  
341 improve the nutritional intake of the diets of young people.

342 The limitations of our study warrant consideration. The size of the cohort subsample is relatively  
343 small and we did not have statistical power to investigate longitudinal patterns by ethnic group  
344 however, it is the first cohort study of young adults to specifically focus on ethnic determinants of  
345 health and chronic disease risk. Furthermore, our study contributes to the very limited longitudinal  
346 data on the role of adolescent diet on cardiometabolic risk in young adulthood and is the first such  
347 study in the UK. We achieved a response rate of 97% in our follow-up, which is remarkably high;  
348 this was achieved mainly through the use of social media, which allowed us to locate participants  
349 who had moved from their childhood residence. Our data collection focused on a relatively small  
350 number of cardiometabolic risk factors, which prevented us from assessing all aspects of risk, for  
351 example, the prevalence of Metabolic Syndrome, and making comparisons with the literature in this  
352 regard. Although our study is the first longitudinal study of adolescents and young adults to collect  
353 nutritional intake data, we were limited in our dietary data collection to assessing a single 24-hour  
354 period and therefore we have not assessed micronutrient intakes; this has limited our ability to  
355 investigate the role of, for example vitamin D, in determining cardiometabolic risk. Under-reporting  
356 is a common issue in dietary data collection; we estimated under-reporting to affect 36% of our  
357 recalls, which is comparable to other studies<sup>(36)</sup>. Importantly, under-reporting was not associated with  
358 gender, ethnicity or educational attainment, suggesting there is no systematic bias in our dietary data.  
359 We did not exclude under-reporters from our analyses; our reporting period was relatively short, and  
360 we deemed low energy intakes to be plausible over a 24-hour period. In our dietary behaviours  
361 questionnaire, which asked about daily portions of fruit and vegetables, we provided portion size  
362 guidance to aid the participants in estimating their daily portions, however our guidance was not  
363 exhaustive and did not focus on composite dishes such as soup, which may have led to inaccuracies  
364 in reporting. We did not collect data on fizzy drink consumption at 21-23 years due to time restrictions  
365 for our data collection visits; sugar sweetened beverages are receiving significant attention at present  
366 due to their links with cardiometabolic risk; our data has not enabled us to look at longitudinal patterns  
367 of consumption. Our dietary analysis is also limited to focusing on single nutrients and food groups  
368 and does not consider interactions between these or reflect the complexity of dietary intake; dietary  
369 pattern analysis would enable a better understanding of dietary interactions.

370 In conclusion, findings from this cohort of young adults representing the major ethnic groups of the  
371 UK population demonstrate the detrimental effects of adolescent dietary behaviours, particularly  
372 skipping breakfast on cardiometabolic risk factors. The results also show these behaviours to be  
373 particularly prevalent amongst Black African and Caribbean ethnic groups. Our study highlights the



374 importance of targeting health promotion interventions to improve dietary behaviours such as  
375 breakfast consumption at specific groups of young adults in the population.

## 376 **ACKNOWLEDGEMENTS**

377 We acknowledge the invaluable support of participants and their parents, the Participant Advisory  
 378 Group, schools, civic leaders, local GP surgeries and community pharmacies, the Clinical Research  
 379 Centre at Queen Mary University of London, the Clinical Research Facility at University College  
 380 Hospital, the survey assistants and nurses involved with data collection, Rachel Bonner, senior  
 381 respiratory physiologist (UCL GOS ICH) for spirometry training and over-reading of data for quality  
 382 control, the Primary Care Research Network, and Professor Sanders and Cruickshank at the Diabetes  
 383 and Nutritional Sciences Division at Kings College London for hosting the feasibility study.  
 384 Seeromanie Harding is the Principal Investigator of DASH. All authors contributed to study design,  
 385 analyses and writing of the article. The study was funded by the MRC (MC\_U130015185/MC\_  
 386 UU\_12017/13), Chief Scientist Office (SPHSU13), North Central London Research Consortium and  
 387 the Primary Care Research Network.

388 **Data sharing:** The DASH data are available to researchers via a data request to the MRC Social and  
 389 Public Health Science Unit. Applications and the data sharing policy for DASH can be found at  
 390 [http://dash.sphsu.mrc.  
 391 ac.uk/DASH\\_dsp\\_v1\\_November-2012\\_draft.pdf](http://dash.sphsu.mrc.ac.uk/DASH_dsp_v1_November-2012_draft.pdf). It reflects the MRC  
 392 guidance on data sharing with the aim of making the data as widely and freely available as possible  
 393 while safeguarding the privacy of participants, protecting confidential data, and maintaining the  
 394 reputation of the study. All potential collaborators work with a link person, an experienced DASH  
 395 researcher—to support their access to and analysis of the data. The variable-level metadata is  
 396 available from the study team and also via the MRC Data Gateway.

397 **FINANCIAL SUPPORT:** MRC (MC\_U130015185/MC\_ UU\_12017/13), Chief Scientist Office  
 398 (SPHSU13), North Central London Research Consortium and the Primary Care Research Network.

399 **CONFLICTS OF INTEREST:** the authors declare no conflicts of interest.

400 **AUTHORS CONTRIBUTIONS:** SH, MM, LMG designed research; EZE, AC, CB conducted  
 401 research; PH, CB, MJS, ORM analysed data and performed statistical analyses; LMG wrote  
 402 manuscripts and had primary responsibility for final content.

## 403 REFERENCES

- 404 1. Becker E, Boreham R, Chaudhury M *et al.* (2006) *Health Survey for England 2004. The health of minority*  
 405 *ethnic groups*. London: Joint Health Surveys Unit, National Centre for Social Research, Department of  
 406 Epidemiology and Public Health at the Royal Free and University College Medical School.
- 407 2. Donin AS, Nightingale CM, Owen CG *et al.* (2010) Ethnic differences in blood lipids and dietary intake  
 408 between UK children of black African, black Caribbean, South Asian, and white European origin: the Child  
 409 Heart and Health Study in England (CHASE). *Am J Clin Nutr* **92**, 776-783.
- 410 3. Ogden CL, Flegal KM, Carroll MD *et al.* (2002) Prevalence and trends in overweight among US children  
 411 and adolescents, 1999-2000. *JAMA* **288**, 1728-1732.
- 412 4. Nelson TF, Gortmaker SL, Subramanian SV *et al.* (2007) Disparities in overweight and obesity among US  
 413 college students. *Am J Health Behav* **31**, 363-373.
- 414 5. Zhang Q, Wang Y, Huang ES (2009) Changes in racial/ethnic disparities in the prevalence of Type 2  
 415 diabetes by obesity level among US adults. *Ethn Health* **14**, 439-457.
- 416 6. Donin AS, Nightingale CM, Owen CG *et al.* (2010) Nutritional composition of the diets of South Asian,  
 417 black African-Caribbean and white European children in the United Kingdom: the Child Heart and Health  
 418 Study in England (CHASE). *Br J Nutr* **104**, 276-285.
- 419 7. Harding S, Teyhan A, Maynard MJ *et al.* (2008) Ethnic differences in overweight and obesity in early  
 420 adolescence in the MRC DASH study: the role of adolescent and parental lifestyle. *Int J Epidemiol* **37**, 162-  
 421 172.
- 422 8. Nicklas TA, Myers L, Reger C *et al.* (1998) Impact of breakfast consumption on nutritional adequacy of the  
 423 diets of young adults in Bogalusa, Louisiana: ethnic and gender contrasts. *J Am Diet Assoc* **98**, 1432-1438.
- 424 9. Ritterman Weintraub ML, Fernald LC, Adler N *et al.* (2015) Perceptions of social mobility: development of  
 425 a new psychosocial indicator associated with adolescent risk behaviors. *Front Public Health* **3**, 62.
- 426 10. Donin AS, Nightingale CM, Owen CG *et al.* (2014) Regular breakfast consumption and type 2 diabetes  
 427 risk markers in 9- to 10-year-old children in the child heart and health study in England (CHASE): a cross-  
 428 sectional analysis. *PLoS Med* **11**, e1001703.
- 429 11. Smith KJ, Gall SL, McNaughton SA *et al.* (2010) Skipping breakfast: longitudinal associations with  
 430 cardiometabolic risk factors in the Childhood Determinants of Adult Health Study. *Am J Clin Nutr* **92**, 1316-  
 431 1325.
- 432 12. Berkey CS, Rockett HR, Gillman MW *et al.* (2003) Longitudinal study of skipping breakfast and weight  
 433 change in adolescents. *Int J Obes Relat Metab Disord* **27**, 1258-1266.
- 434 13. Szajewska H, Rusczyński M (2010) Systematic review demonstrating that breakfast consumption  
 435 influences body weight outcomes in children and adolescents in Europe. *Crit Rev Food Sci Nutr* **50**, 113-119.
- 436 14. Merten MJ, Williams AL, Shriver LH (2009) Breakfast consumption in adolescence and young adulthood:  
 437 parental presence, community context, and obesity. *J Am Diet Assoc* **109**, 1384-1391.
- 438 15. Watts AW, Mason SM, Loth K *et al.* (2016) Socioeconomic differences in overweight and weight-related  
 439 behaviors across adolescence and young adulthood: 10-year longitudinal findings from Project EAT. *Prev*  
 440 *Med* **87**, 194-199.
- 441 16. Harding S, Whitrow M, Maynard MJ *et al.* (2007) Cohort profile: The DASH (Determinants of Adolescent  
 442 Social well-being and Health) Study, an ethnically diverse cohort. *Int J Epidemiol* **36**, 512-517.
- 443 17. Nelson M, Holmes B, Roberts C *et al.* (2013) *Dietary assessment - Recalls. Diet and physical activity*  
 444 *measurement toolkit*. <http://dapa-toolkit.mrc.ac.uk/dietary-assessment/methods/recalls/index.html>: MRC.
- 445 18. Nelson, Atkinson, Meyer J (1997) *A photographic atlas of food portion sizes*: MAFF Publications, London,  
 446 UK.
- 447 19. Goff LM, Timbers L, Style H *et al.* (2014) Dietary intake in Black British adults; an observational  
 448 assessment of nutritional composition and the role of traditional foods in UK Caribbean and West African  
 449 diets. *Public Health Nutr*, 1-11.
- 450 20. Stadlmayr B, Charrondiere RVE, Bayli R *et al.* (2012) *West African Food Composition Table*. Rome.
- 451 21. Department of Health (2011) National Diet and Nutrition Survey: Headline results from Years 1 and 2  
 452 (combined) of the rolling programme 2008-9 - 2009-10. England: Department of Health.
- 453 22. Henry CJ (2005) Basal metabolic rate studies in humans: measurement and development of new  
 454 equations. *Public Health Nutr* **8**, 1133-1152.

23. Rennie KL, Coward A, Jebb SA (2007) Estimating under-reporting of energy intake in dietary surveys using an individualised method. *Br J Nutr* **97**, 1169-1176.
24. Dowd KP, Harrington DM, Donnelly AE (2012) Criterion and concurrent validity of the activPAL professional physical activity monitor in adolescent females. *PLoS One* **7**, e47633.
25. Gower BA, Nagy TR, Goran MI (1999) Visceral fat, insulin sensitivity, and lipids in prepubertal children. *Diabetes* **48**, 1515-1521.
26. Ramachandran A, Snehalatha C, Viswanathan V *et al.* (1997) Risk of noninsulin dependent diabetes mellitus conferred by obesity and central adiposity in different ethnic groups: a comparative analysis between Asian Indians, Mexican Americans and Whites. *Diabetes Res Clin Pract* **36**, 121-125.
27. Herman WH, Ma Y, Uwaifo G *et al.* (2007) Differences in A1C by race and ethnicity among patients with impaired glucose tolerance in the Diabetes Prevention Program. *Diabetes Care* **30**, 2453-2457.
28. Ziemer DC, Kolm P, Weintraub WS *et al.* (2010) Glucose-independent, black-white differences in hemoglobin A1c levels: a cross-sectional analysis of 2 studies. *Ann Intern Med* **152**, 770-777.
29. Vyas A, Greenhalgh A, Cade J *et al.* (2003) Nutrient intakes of an adult Pakistani, European and African-Caribbean community in inner city Britain. *J Hum Nutr Diet* **16**, 327-337.
30. Goff LM, Griffin BA, Lovegrove JA *et al.* (2013) Ethnic differences in beta-cell function, dietary intake and expression of the metabolic syndrome among UK adults of South Asian, black African-Caribbean and white-European origin at high risk of metabolic syndrome. *Diab Vasc Dis Res* **10**, 315-323.
31. Miller GJ, Kotecha S, Wilkinson WH *et al.* (1988) Dietary and other characteristics relevant for coronary heart disease in men of Indian, West Indian and European descent in London. *Atherosclerosis* **70**, 63-72.
32. McKeigue PM, Marmot MG, Adelstein AM *et al.* (1985) Diet and risk factors for coronary heart disease in Asians in northwest London. *Lancet* **2**, 1086-1090.
33. Knight TM, Smith Z, Whittles A *et al.* (1992) Insulin resistance, diabetes, and risk markers for ischaemic heart disease in Asian men and non-Asian in Bradford. *Br Heart J* **67**, 343-350.
34. Sevak L, McKeigue PM, Marmot MG (1994) Relationship of hyperinsulinemia to dietary intake in south Asian and European men. *Am J Clin Nutr* **59**, 1069-1074.
35. Deshmukh-Taskar P, Nicklas TA, Radcliffe JD *et al.* (2013) The relationship of breakfast skipping and type of breakfast consumed with overweight/obesity, abdominal obesity, other cardiometabolic risk factors and the metabolic syndrome in young adults. The National Health and Nutrition Examination Survey (NHANES): 1999-2006. *Public Health Nutr* **16**, 2073-2082.
36. Poslusna K, Ruprich J, de Vries JH *et al.* (2009) Misreporting of energy and micronutrient intake estimated by food records and 24 hour recalls, control and adjustment methods in practice. *Br J Nutr* **101 Suppl 2**, S73-85.

**Table 1. Dietary intake, anthropometry and cardiometabolic risk factors of participants, at 21-23y, in the Determinants of Adolescent, now young Adults, Social well-being and Health study**

	<b>Whole cohort (n=665)</b>	<b>White British (n=107)</b>	<b>Black Caribbean (n=102)</b>	<b>Black African (n=132)</b>	<b>Indian (n=98)</b>	<b>Pakistani Bangladeshi (n=111)</b>	<b>Other (n=115)</b>	
	<b>Mean (95% CI)</b>	<b>Mean (95% CI)</b>	<b>Mean (95% CI)</b>	<b>Mean (95% CI)</b>	<b>Mean (95% CI)</b>	<b>Mean (95% CI)</b>	<b>Mean (95% CI)</b>	<b><i>p</i></b>
<b>Male/Female (n)</b>	329/336	52/55	47/55	58/74	52/46	62/49	57/57	0.842
<b>Age (y)</b>	22.7 (22.7, 22.8)	22.7 (22.5, 22.8)	22.7 (22.5, 22.9)	22.6 (22.4, 22.9)	22.9 (22.7, 23.1)	22.6 (22.7, 23.1)	22.8 (22.6, 23.0)	0.129
<b>Weight (kg)</b>	71.6 (70.0, 73.3)	72.5 (68.8, 76.2)	76.6 <sup>a</sup> (71.7, 81.5)	77.1 <sup>b</sup> (73.0, 81.2)	66.3 <sup>a,b</sup> (62.7, 69.9)	68.0 <sup>a,b</sup> (64.1, 71.9)	69.9 (66.5, 73.3)	<0.001
<b>Height (cm)</b>	170 (169, 171)	172 <sup>a</sup> (170, 175)	170 (168, 173)	172 <sup>b</sup> (170, 174)	168 <sup>a,b</sup> (166, 171)	168 <sup>a,b</sup> (165, 170)	169 (167, 171)	<0.001
<b>BMI (kg/m<sup>2</sup>)</b>	24.7 (24.2, 25.2)	24.2 <sup>a,c</sup> (23.1, 25.4)	26.2 <sup>a,b</sup> (24.8, 27.7)	26.1 <sup>c,d</sup> (24.8, 27.4)	23.4 <sup>b,d</sup> (22.2, 24.5)	24.0 (22.9, 25.2)	24.4 (23.4, 25.5)	<0.001
<b>Overweight/obese (%)<sup>a</sup></b>	39	35	50	44	36	36	35	0.063
<b>Waist (cm)</b>	82.2 (80.9, 83.5)	83.0 (80.2, 85.8)	83.6 (80.0, 87.3)	83.0 (79.9, 86.1)	80.8 (77.6, 84.1)	81.7 (79.0, 84.4)	81.2 (78.4, 84.1)	0.878
<b>Waist:height ratio</b>	0.48 (0.48, 0.49)	0.48 (0.46, 0.50)	0.49 (0.47, 0.51)	0.48 (0.46, 0.50)	0.48 (0.46, 0.50)	0.49 (0.47, 0.50)	0.48 (0.46, 0.50)	0.429
<b>Total cholesterol (mmol/l)</b>	4.35 (4.26, 4.43)	4.34 (4.17, 4.52)	4.34 (4.14, 4.54)	4.07 <sup>a</sup> (3.93, 4.22)	4.36 (4.21, 4.51)	4.53 <sup>a</sup> (4.35, 4.72)	4.33 (4.14, 4.52)	0.007
<b>HDL cholesterol (mmol/l)</b>	1.54 (1.50, 1.57)	1.56 (1.49, 1.64)	1.61 <sup>a</sup> (1.53, 1.70)	1.60 <sup>b</sup> (1.52, 1.67)	1.41 <sup>a,b</sup> (1.34, 1.48)	1.47 (1.40, 1.54)	1.55 (1.46, 1.64)	0.003
<b>HbA1c (mmol/mol)</b>	34 (34, 35)	34 (33, 35)	35 (34, 36)	35 (34, 36)	35 (34, 36)	34 (33, 35)	34 (33, 35)	0.164
<b>Systolic blood pressure (mmHg)</b>	114 (113, 115)	117 (114, 120)	116 (113, 119)	116 (113, 119)	113 (110, 116)	112 (109, 115)	113 (110, 115)	0.090
<b>Diastolic blood pressure (mmHg)</b>	72 (72, 73)	73 (71, 75)	74 (72, 76)	74 (72, 76)	72 (70, 74)	72 (70, 74)	70 (68, 72)	0.073
<b>Energy intake (kcal/day)</b>	2192 (2097, 2288)	2008 (1811, 2205)	2242 (2018, 2466)	1976 (1766, 2186)	2423 (2164, 2682)	2260 (2001, 2520)	2238 (1982, 2494)	0.255

<b>Fat intake</b>	38.0	37.9	38.5	36.5	35.6	41.1	38.6	0.168
(% of energy)	(37.0, 39.0)	(35.5, 40.2)	(36.4, 40.5)	(33.7, 39.3)	(33.0, 38.3)	(38.9, 43.4)	(36.1, 41.1)	
<b>Saturated fat intake</b>	14.4	15.5	13.9	13.0 <sup>a</sup>	13.1 <sup>a</sup>	16.4 <sup>a</sup>	14.5	0.031
(% of energy)	(13.9, 15.0)	(14.2, 16.9)	(12.7, 15.2)	(11.7, 14.3)	(11.8, 14.4)	(14.9, 17.9)	(13.3, 15.6)	
<b>Carbohydrate intake</b>	47.4	46.1	48.1	47.9	50.4	45.6	46.5	0.170
(% of energy)	(46.3, 48.5)	(43.8, 48.4)	(45.4, 50.7)	(44.8, 51.0)	(47.8, 53.0)	(42.9, 48.3)	(43.8, 49.2)	
<b>Protein intake</b>	16.4	16.0	15.9	17.1	15.7	16.6	16.8	0.953
(% of energy)	(15.7, 17.1)	(14.4, 17.6)	(14.2, 17.6)	(14.8, 19.5)	(13.9, 17.4)	(15.1, 18.1)	(15.6, 18.2)	
<b>Sugar intake</b>	17.1	16.6	19.5	18.8	16.2	14.5	17.1	0.151
(% of energy)	(16.2, 18.0)	(14.5, 18.8)	(17.0, 22.0)	(16.1, 21.6)	(14.2, 18.2)	(12.7, 16.2)	(15.0, 19.3)	
<b>Fibre intake</b>	8.1	9.3 <sup>a</sup>	7.3 <sup>a,c</sup>	7.7 <sup>b,c</sup>	8.8 <sup>b,c</sup>	7.3 <sup>c</sup>	8.0	<0.001
(g/1000kcal)	(7.7, 8.5)	(8.1, 10.5)	(6.3, 8.2)	(6.8, 8.7)	(7.6, 9.9)	(6.3, 8.3)	(6.9, 9.0)	
<b>Sodium intake</b>	1285	1407	1320	1122	1192	1314	1345	0.082
(mg/1000kcal)	(1231, 1340)	(1270, 1544)	(1157, 1483)	(1010, 1235)	(1054, 1330)	(1191, 1436)	(1218, 1471)	
<b>Sedentary time</b>	738	798	670	833	772	720	692	0.234
(mins/day)	(683, 792)	(650, 947)	(573, 767)	(6565, 1000)	(654, 889)	(560, 880)	(575, 807)	
<b>Skipping breakfast (%)</b>	56	53	66	64	46	51	54	0.118
<b>Low fruit consumption (%)</b>	57	52	59	65	55	56	52	0.373
<b>Low vegetable consumption (%)</b>	63	54	75	73	55	73*	51*	<0.001
<b>Smoker (%)</b>	24	47*	21	11*	24	24	24	0.001
<b>Educated to degree level (%)</b>	47	40	34*	58*	57	49	44	0.001

BMI, body mass index. Overweight/obese defined as BMI>25kg/m<sup>2</sup>. Low fruit consumption defined as less than 2 portions per day; low vegetable consumption defined as less than 3 portions per day; skipping breakfast defined as not eating breakfast every day of the week. Continuous variables are shown as mean (95% CI), categorical variables shown as n or %, as indicated in the variable label. Differences between ethnic groups in continuous variables were tested with one-way ANOVA with post-hoc analysis, the same superscript letters within a row denote significant differences between those ethnic groups; ethnic differences in frequency variables were tested with chi-square test with post-hoc analysis, differences are denoted with \*.

**Table 2. Influence of dietary behaviours at 11-13 years and at 21-23years on cardiometabolic risk factors at 21-23 years. The Determinants of Adolescent, now young Adults, Social well-being and Health study.**

	BMI		Waist:height ratio		Total cholesterol		HDL-cholesterol		HbA1c		Systolic blood pressure	
<b>11-13y</b>	Coef. (95% CI)	<i>p</i>	Coef. (95% CI)	<i>p</i>	Coef. (95% CI)	<i>p</i>	Coef. (95% CI)	<i>p</i>	Coef. (95% CI)	<i>p</i>	Coef. (95% CI)	<i>p</i>
Skipping breakfast (No - Ref)												
Yes	1.41 (0.57, 2.26)	<0.001	0.02 (0.01, 0.03)	<0.001	0.15 (-0.01, 0.31)	0.063	0.01 (-0.06, 0.08)	0.725	0.43 (-0.53, 1.39)	0.379	0.74 (-0.89, 2.38)	0.370
Not stated	0.78 (-0.70, 2.26)	0.302	0.01 (-0.01, 0.03)	0.353	0.16 (-0.12, 0.45)	0.266	0.02 (-0.12, 0.15)	0.820	0.05 (-1.66, 1.75)	0.958	0.33 (-2.54, 3.19)	0.823
Fruit ( $\geq 2$ serves/day - Ref)												
<2 serves/day	0.68 (-0.17, 1.53)	0.115	0.01 (0.00, 0.03)	0.034	-0.004 (-0.16, 0.16)	0.958	-0.05 (-0.12, 0.02)	0.197	-0.16 (-1.11, 0.79)	0.742	0.66 (-0.97, 2.30)	0.425
Not stated	0.65 (-0.58, 1.90)	0.299	0.01 (-0.01, 0.03)	0.225	0.03 (-0.20, 0.27)	0.779	-0.003 (-0.11, 0.10)	0.951	-0.70 (-2.10, 0.70)	0.325	0.21 (-2.17, 2.59)	0.865
Vegetables ( $\geq 3$ serves/day - Ref)												
<3 serves/day	-0.03 (-0.89, 0.82)	0.941	0.002 (-0.01, 0.01)	0.723	-0.17 (-0.33, -0.01)	0.033	-0.007 (-0.06, 0.08)	0.838	0.18 (-0.75, 1.12)	0.702	-0.003 (-1.64, 1.63)	0.997
Not stated	-1.70 (-1.00, 1.29)	0.788	-0.004 (-0.03, 0.02)	0.709	-0.11 (-0.40, 0.17)	0.427	-0.007 (-0.12, 0.14)	0.905	-0.38 (-2.07, 1.30)	0.656	-0.79 (-3.66, 2.09)	0.591
Fizzy drinks (No - Ref)												
Yes	0.03 (-0.89, 0.84)	0.947	0.001 (-0.01, 0.01)	0.888	0.05 (-0.11, 0.21)	0.542	-0.04 (-0.11, 0.03)	0.240	-0.34 (-1.30, 0.62)	0.488	1.77 (0.12, 3.43)	0.036
Not stated	0.06 (-1.44, 1.56)	0.938	-0.002 (-0.02, 0.02)	0.858	0.05 (-0.24, 0.34)	0.742	-0.007 (-0.14, 0.12)	0.920	-0.53 (-2.25, 1.19)	0.546	1.22 (-1.64, 4.09)	0.403
<b>21-23y</b>												
Skipping breakfast (No - Ref)												
Yes	1.05 (0.22, 1.89)	0.014	0.02 (0.01, 0.03)	0.002	0.22 (0.06, 0.37)	0.007	0.0002 (-0.07, 0.07)	0.995	0.73 (-0.22, 1.67)	0.130	0.44 (-1.18, 2.05)	0.597
Not stated	-1.32 (-4.44, 1.80)	0.406	-0.01 (-0.05, 0.04)	0.731	0.30 (-0.26, 0.85)	0.292	0.07 (-0.17, 0.32)	0.556	0.12 (-2.93, 3.18)	0.937	-3.56 (-9.56, 2.44)	0.244
Fruit ( $\geq 2$ serves/day - Ref)												
<2 serves/day	-0.26 (-1.11, 0.59)	0.552	0.001 (-0.01, 0.01)	0.912	-0.12 (-0.28, 0.04)	0.147	-0.04 (-0.11, 0.03)	0.270	-0.50 (-1.46, 0.46)	0.305	0.26 (-1.37, 1.88)	0.758
Not stated	-1.65 (-4.32, 1.02)	0.225	-0.008 (-0.05, 0.03)	0.676	0.25 (-0.23, 0.74)	0.301	0.03 (-0.19, 0.24)	0.809	-0.11 (-2.95, 2.74)	0.940	-0.39 (-5.51, 4.73)	0.881

Vegetables ( $\geq 3$ serves/day - Ref)												
<3 serves/day	-0.07	0.873	0.001	0.859	-0.06	0.456	-0.02	0.630	0.29	0.571	-0.72	0.420
	(-0.99, 0.84)		(-0.01, 0.01)		(-0.24, 0.11)		(-0.09, 0.06)		(-0.72, 1.30)		(-2.49, 1.04)	
Not stated	-1.40	0.221	-0.01	0.574	-0.12	0.559	0.08	0.385	-0.17	0.884	-3.27	0.137
	(-3.66, 0.85)		(-0.04, 0.02)		(-0.51, 0.28)		(-0.10, 0.25)		(-2.49, 2.15)		(-7.58, 1.05)	

BMI, body mass index. Skipping breakfast defined as eating breakfast less than 5 days per week; low fruit consumption defined as less than 2 portions per day; low vegetable consumption defined as less than 3 portions per day; fizzy drink consumption defined as consuming fizzy drinks every day or most days. Multivariate linear regression models for BMI, waist:height ratio, total cholesterol, HDL-cholesterol, HbA1c and systolic blood pressure (separate models); coefficients were estimated with linear regression, adjusted for gender and ethnicity at 11-13 years, and additionally adjusted for educational attainment (attainment/non-attainment of higher education degree), physical activity (based on tertiles of sitting/sedentary time), smoking (current smoker/non-smoker), energy intake and saturated fat intake at 21-23 years. N=664, 635, 499, 484, 481, and 664, respectively, for linear regression models with BMI, waist:height ratio, total cholesterol, HDL-cholesterol, HbA1c, and systolic blood pressure, respectively, as the outcome.



**Table 3. Influence of dietary behaviours at 11-13 years and at 21-23years on nutritional composition of the diet at 21-23years. The Determinants of Adolescent, now young Adults, Social well-being and Health study.**

	Energy intake (kcal)		Fat (% of energy)		Saturated fat (% of energy)		Sugar (% of energy)		Fibre (g/1000kcal)		Sodium (mg/1000kcal)	
<b>11-13y</b>	Coef. (95% CI)	<i>p</i>	Coef. (95% CI)	<i>p</i>	Coef. (95% CI)	<i>p</i>	Coef. (95% CI)	<i>p</i>	Coef. (95% CI)	<i>p</i>	Coef. (95% CI)	<i>p</i>
Skipping breakfast (No - Ref)												
Yes	-116.25 (-263.65, 31.15)	0.122	-0.19 (-1.86, 1.48)	0.826	0.42 (-0.48, 1.31)	0.361	0.11 (-1.43, 1.65)	0.889	-0.81 (-1.49, -0.12)	0.022	-58.83 (-184.66, 67.00)	0.359
Not stated	-176.37 (-435.51, 82.77)	0.182	0.14 (-2.80, 3.08)	0.927	0.66 (-0.92, 2.24)	0.411	2.13 (-0.58, 4.85)	0.123	-0.74 (-1.95, 0.47)	0.230	50.38 (-170.84, 271.61)	0.655
Fruit (≥2 serves/day - Ref)												
<2 serves/day	142.89 (-4.17, 289.95)	0.057	-0.57 (-2.24, 1.10)	0.503	-0.44 (-1.33, 0.46)	0.339	-0.14 (-1.68, 1.40)	0.859	-0.36 (-1.04, 0.33)	0.312	-5.49 (-131.17, 120.19)	0.932
Not stated	23.48 (-191.86, 238.83)	0.830	-0.84 (-3.29, 1.60)	0.498	-0.65 (-1.96, 0.66)	0.330	1.34 (-0.92, 3.59)	0.244	0.16 (-0.85, 1.17)	0.752	68.67 (-115.37, 252.70)	0.464
Vegetables (≥3 serves/day - Ref)												
<3 serves/day	-48.49 (-197.09, 100.11)	0.522	-1.15 (-2.83, 0.53)	0.178	-0.60 (-1.50, 0.30)	0.193	1.32 (-0.23, 2.86)	0.095	-0.31 (-1.00, 0.39)	0.383	-27.77 (-154.45, 98.91)	0.667
Not stated	-108.10 (-368.68, 152.47)	0.416	-1.23 (-4.17, 1.72)	0.414	-0.07 (-1.65, 1.51)	0.930	3.56 (0.85, 6.27)	0.010	-0.10 (-1.32, 1.12)	0.869	47.31 (-174.83, 269.45)	0.676
Fizzy drinks (No - Ref)												
Yes	34.82 (-115.70, 185.35)	0.650	0.04 (-1.66, 1.74)	0.964	0.34 (-0.58, 1.25)	0.469	-0.08 (-1.65, 1.48)	0.915	-0.72 (-1.42, -0.02)	0.045	-51.30 (-179.49, 76.89)	0.432
Not stated	-47.59 (-308.79, 213.60)	0.721	-0.27 (-3.23, 2.69)	0.857	0.66 (-0.93, 2.25)	0.414	3.15 (0.44, 5.87)	0.023	-0.66 (-1.88, 0.56)	0.287	70.21 (-152.23, 292.64)	0.536
<b>21-23y</b>												
Skipping breakfast (No - Ref)												
Yes	-68.16 (-214.82, 78.49)	0.362	1.10 (-0.56, 2.75)	0.195	0.26 (-0.63, 1.15)	0.565	-0.49 (-2.02, 1.04)	0.530	-1.33 (-2.01, -0.65)	<0.001	-35.13 (-160.14, 89.88)	0.581
Not stated	-144.67 (-689.91, 400.57)	0.602	1.70 (-4.46, 7.87)	0.587	-0.99 (-4.30, 2.33)	0.558	-2.17 (-7.87, 3.54)	0.456	0.00 (-2.52, 2.52)	0.998	-199.83 (-664.60, 264.93)	0.399
Fruit (≥2 serves/day - Ref)												
<2 serves/day	27.47 (-119.98, 174.91)	0.715	0.70 (-0.97, 2.37)	0.411	0.89 (-0.01, 1.78)	0.052	-0.47 (-2.01, 1.07)	0.547	-1.22 (-1.90, -0.54)	<0.001	-29.32 (-154.98, 96.35)	0.647
Not stated	-7.71 (-473.24, 457.83)	0.974	1.19 (-4.08, 6.45)	0.658	0.14 (-2.68, 2.96)	0.921	-3.14 (-8.00, 1.72)	0.205	0.67 (-1.48, 2.82)	0.542	-33.96 (-430.73, 362.82)	0.867

Vegetables ( $\geq 3$ serves/day - Ref)												
<3 serves/day	-27.94	0.731	1.50	0.100	1.06	0.032	0.11	0.896	-1.30	0.001	-101.31	0.143
	(-187.19, 131.32)		(-0.29, 3.30)		(0.09, 2.02)		(-1.55, 1.78)		(-2.04, -0.56)		(-236.80, 34.18)	
Not stated	26.10	0.896	4.01	0.076	1.27	0.294	-1.83	0.380	-0.76	0.410	18.78	0.912
	(-366.17, 418.37)		(-0.42, 8.43)		(-1.10, 3.65)		(-5.93, 2.27)		(-2.58, 1.05)		(-314.96, 352.51)	

Skipping breakfast defined as eating breakfast less than 5 days per week; low fruit consumption defined as less than 2 portions per day; low vegetable consumption defined as less than 3 portions per day; fizzy drink consumption defined as consuming fizzy drinks every day or most days. Multivariate linear regression models for energy intake, % energy from fat, % energy from saturated fat, % energy from sugar, fibre intake, sodium intake (separate models); coefficients were estimated with linear regression, adjusted for gender, and ethnicity at 11-13 years, and additionally adjusted for educational attainment (attainment/non-attainment of higher education degree), physical activity (based on tertiles of sitting/sedentary time), and smoking (current smoker/non-smoker) at 21-23 years. N=628 for linear regression models for all six nutritional composition variables as the outcome.

**FIGURE LEGENDS**

**Figure 1.** Rates of (a) regularly skipping breakfast, (b) low daily fruit and (c) low daily vegetable consumption according to ethnicity at 11-13 years and at 21-23 years. The Determinants of Adolescent, now young Adults, Social well-being and Health study.



